

标题: Probing the mass and structure of the Ring Nebula in Lyra with SOFIA/GREAT observations of the [CII] 158 micron line

作者: Sahai, R (Sahai, R.); Morris, MR (Morris, M. R.); Werner, MW (Werner, M. W.); Guesten, R (Guesten, R.); Wiesemeyer, H (Wiesemeyer, H.); Sandell, G (Sandell, G.)

来源出版物: ASTRONOMY & ASTROPHYSICS 卷: 542 文献号: L20 DOI: 10.1051/0004-6361/201219021 出版年: JUN 2012

在 Web of Science 中的被引频次: 0

被引频次合计: 0

引用的参考文献数: 20

摘要: We have obtained new velocity-resolved spectra of the [CII] 158  $\mu\text{m}$  line towards the Ring Nebula in Lyra (NGC 6720), one of the best-studied planetary nebulae, in order to probe its controversial 3-dimensional structure and to estimate the mass of circumstellar material in this object. We used the terahertz receiver GREAT aboard the SOFIA airborne telescope to obtain the [CII] spectra at eight locations within and outside the bright optical ring of NGC 6720. Emission was detected at all positions except for the most distant position along the nebula's minor axis, and generally covers a broad velocity range,  $\Delta V$  similar to 50  $\text{km s}^{-1}$  (FWZI), except at a position along the major axis located just outside the optical ring, where it is significantly narrower ( $\Delta V$  similar to 25  $\text{km s}^{-1}$ ). The one narrow spectrum appears to be probing circumstellar material lying outside the main nebular shell that has not been accelerated by past fast wind episodes from the central star, and therefore most likely comes from equatorial and/or low-latitude regions of this multipolar nebula. Along lines-of-sight passing within about 10" of the nebular center, the CII column density is a factor 46 higher than the CO column density. The total mass of gas associated with the [CII] emission inside a circular region of diameter 87".5 is at least 0.11  $M_{\odot}$ . A significant amount of [CII] flux arises from a photodissociation region immediately outside the bright optical ring, where we find a CII to CO ratio of  $>6.5$ , lower than that seen towards the central region. Comparing our data with lower-quality CI spectra, which indicate similarly large CI/CO ratios in NGC 6720, we conclude that the bulk of elemental carbon in NGC 6720 is divided roughly equally between CII and CI, and that the emissions from these species are far more robust tracers of circumstellar material than CO in this object and other evolved planetary nebulae.

入藏号: WOS:000305803300128

语种 : English

文献类型: Article

作者关键词: planetary nebulae: individual: NGC 6720; stars: winds, outflows; photon-dominated region (PDR); circumstellar matter; planetary nebulae: general

地址: [Sahai, R.; Werner, M. W.] CALTECH, Jet Prop Lab, Pasadena, CA 91109 USA

[Morris, M. R.] Univ Calif Los Angeles, Dept Phys & Astrophys, Los Angeles, CA 90095 USA

[Guesten, R.; Wiesemeyer, H.] Max Planck Inst Radioastron, D-53121 Bonn, Germany

[Sandell, G.] NASA, SOFIA USRA, Ames Res Ctr, Moffett Field, CA 94035 USA

通讯作者地址: Sahai, R (通讯作者), CALTECH, Jet Prop Lab, 4800 Oak Grove Dr, Pasadena, CA 91109 USA.

电子邮件地址: raghvendra.sahai@jpl.nasa.gov

出版商: EDP SCIENCES S A

出版商地址: 17, AVE DU HOGGAR, PA COURTABOEUF, BP 112, F-91944 LES ULIS CEDEX  
A, FRANCE

Web of Science 类别: Astronomy & Astrophysics

研究方向: Astronomy & Astrophysics

IDS 号: 965YI

ISSN: 0004-6361

29 字符的来源出版物名称缩写: ASTRON ASTROPHYS

ISO 来源出版物缩写: Astron. Astrophys.

来源出版物页码计数: 4